

DESY 02-046

April 2002

Exceptional Coset Spaces and Unification in Six Dimensions

T. Asaka, W. Buchmüller, L. Covi

Deutsches Elektronen-Synchrotron DESY, Hamburg, Germany

Abstract

The coset spaces $E_8/SO(10) \times H_F$ allow complex structures which can account for three quark-lepton generations including right-handed neutrinos. We show that in the context of supersymmetric $SO(10)$ gauge theories in 6 dimensions they also provide the Higgs fields which are needed to break the electroweak and $B - L$ gauge symmetries, and to generate small neutrino masses via the seesaw mechanism.

The standard model gauge groups of electroweak and strong interactions are naturally unified in the simple gauge group $SU(5)$ [1]. With the increasing experimental evidence for neutrino masses and mixings the larger gauge group $SO(10)$ [2] with the Pati-Salam subgroup $SU(4) \times SU(2) \times SU(2)$ [3] appears particularly attractive, since all quarks and leptons of a single generation, including the right-handed neutrino, are then also unified in a single multiplet.

The unification groups $SU(5)=E_4$, $SO(10)=E_5$ and E_6 [4] belong to the sequence of exceptional groups E_n which terminates at E_8 . This largest exceptional group is attractive for unification [5, 6] since its smallest multiplet, the 248-dimensional adjoint representation, is large enough to accommodate all three generations of quarks and leptons. The theory is naturally supersymmetric. However, in addition to the three known quark-lepton generations the theory also predicts three light mirror generations, contrary to observation.

The group E_8 also appears in the ten-dimensional Yang-Mills supergravity theory where a cancellation of all gauge and gravitational anomalies is obtained by means of the Green-Schwarz mechanism for the group $E_8 \times E_8$ [7]. Compactification to four dimensions [8, 9] can yield low energy effective theories with unbroken $N = 1$ supersymmetry and chiral fermions, similar to the structure of the standard model. The number of families is determined by the Euler characteristic of the compact manifold.

Further, the group E_8 has been considered in attempts to relate quarks and leptons to a coset space G/H , where G is an appropriate simple group and H contains the standard model gauge group [10]. By pairing the scalar degrees of freedom of G/H into complex fields, which become the superpartner of quarks and leptons, the problem of mirror families can be avoided [11]. Particularly attractive are coset spaces $E_8/(SO(10) \times H_F)$ where H_F is a subgroup of $SU(3) \times U(1)$ [11]-[14]. In the case $H_F = SU(3) \times U(1)$ the representation of chiral multiplets is unique,

$$\Omega = (\mathbf{16}, \mathbf{3})_1 + (\mathbf{16}^*, \mathbf{1})_3 + (\mathbf{10}, \mathbf{3}^*)_2 + (\mathbf{1}, \mathbf{3})_4 . \quad (1)$$

Hence, in addition to three quark-lepton generations contained in the three $\mathbf{16}$'s of $SO(10)$, one mirror generation, $\mathbf{16}^*$, occurs, again in contrast to observation. For $H_F = SU(2) \times U(1)^2$ and $H_F = U(1)^3$ various complex structures are possible [13], which contain either three $\mathbf{16}$'s and one $\mathbf{16}^*$, or two $\mathbf{16}$'s and two $\mathbf{16}^*$'s in addition to three $\mathbf{10}$'s and $SO(10)$ singlets.

Orbifold compactifications [9] have recently been applied to GUT field theories. The breaking of the GUT symmetry then automatically yields the required doublet-triplet splitting of Higgs fields [15]. Several $SU(5)$ models have been constructed in 5 dimensions [15]-[18], whereas 6 dimensions are required for the breaking of $SO(10)$ [19, 20]. The $N=2$




Figure 1: Orbifold $T^2/(Z_2^I \times Z_2^{PS} \times Z_2^{GG})$ with the fixpoints O , O_{PS} , O_{GG} , and O_{fl} .

supersymmetric $SO(10)$ theory in 6 dimensions can also be used as a starting point to obtain a $SU(5)$ GUT with three generations [21].

In the following we want to point out that the complex structure (1), when combined with the orbifold compactification of the 6d $SO(10)$ theory, yields in a simple way the supersymmetric standard model with right-handed neutrinos. The three $\mathbf{16}$'s, which contain quarks and leptons, are brane fields; the $\mathbf{16}^*$ and the three $\mathbf{10}$'s are bulk fields. Vacuum expectation values of bulk fields break the symmetries $SU(2) \times U(1)_Y$ and $U(1)_X$, which are left unbroken by the orbifold compactification.

Consider the $SO(10)$ gauge theory in 6d with $N=1$ supersymmetry. The gauge fields $V_M(x, y, z)$, with $M = \mu, 5, 6$, $x^5 = y$, $x^6 = z$, and the gauginos λ_1, λ_2 are conveniently grouped into vector and chiral multiplets of the unbroken $N=1$ supersymmetry in 4d,

$$V = (V_\mu, \lambda_1), \quad \Sigma = (V_{5,6}, \lambda_2). \quad (2)$$

Here V and Σ are matrices in the adjoint representation of $SO(10)$. Symmetry breaking is achieved by compactification on the orbifold $T^2/(Z_2^I \times Z_2^{PS} \times Z_2^{GG})$. The discrete symmetries Z_2 break the extended supersymmetry in 4d; they also break the $SO(10)$ gauge group down to the subgroups $SO(10)$, $G_{PS}=SU(4) \times SU(2) \times SU(2)$ and $G_{GG}=SU(5) \times U(1)_X$, respectively, at three different fixpoints,

$$P_I V(x, -y, -z) P_I^{-1} = \eta_I V(x, y, z) , \quad (3)$$

$$P_{PS} V(x, -y + \pi R_5/2, -z) P_{PS}^{-1} = \eta_{PS} V(x, y + \pi R_5/2, z) , \quad (4)$$

$$P_{GG} V(x, -y, -z + \pi R_6/2) P_{GG}^{-1} = \eta_{GG} V(x, y, z + \pi R_6/2) . \quad (5)$$

Here $P_I = I$, the matrices P_{PS} and P_{GG} are given in ref. [19], and the parities are chosen as $\eta_I = \eta_{PS} = \eta_{GG} = +1$. The extended supersymmetry is broken by choosing in the corresponding equations for Σ all parities $\eta_i = -1$. At the fixpoints in the extra dimensions, $O = (0, 0)$, $O_{PS} = (\pi R_5/2, 0)$ and $O_{GG} = (0, \pi R_6/2)$ the unbroken subgroups are $SO(10)$, G_{PS} and G_{GG} , respectively. In addition, there is a fourth fixpoint at $O_{fl} = (\pi R_5/2, \pi R_6/2)$ [20], which is obtained by combining the three discrete symmetries Z_2 , Z_2^{PS} and Z_2^{GG} defined above,

$$P_{fl} V(x, -y + \pi R_5/2, -z + \pi R_6/2) P_{fl}^{-1} = +V(x, y + \pi R_5/2, z + \pi R_6/2) . \quad (6)$$

The unbroken subgroup at the fixpoint O_{fl} is flipped $SU(5)$, i.e. $G_{fl} = SU(5)' \times U(1)'$. The physical region is obtained by folding the shaded regions in fig. 1 along the dotted line and gluing the edges. The result is a ‘pillow’ with the four fixpoints as corners. The unbroken gauge group of the effective 4d theory is given by the intersection of the $SO(10)$ subgroups at the fixpoints. In this way one obtains the standard model group with an additional $U(1)$ factor, $G_{SM'} = SU(3) \times SU(2) \times U(1)_Y \times U(1)_X$. The difference of baryon and lepton number is the linear combination $B - L = \sqrt{\frac{16}{15}} Y - \sqrt{\frac{8}{5}} X$. The zero modes of the vector multiplet V form the gauge fields of $G_{SM'}$.

The vector multiplet V is a **45**-plet of $SO(10)$ which has an irreducible anomaly in 6 dimensions. It is related to the irreducible anomalies of hypermultiplets in the fundamental and the spinor representations by (cf. [23]),

$$a(\mathbf{45}) = -2a(\mathbf{10}) , \quad a(\mathbf{16}) = a(\mathbf{16}^*) = -a(\mathbf{10}) . \quad (7)$$

Hence, the anomaly of the vector multiplet can be cancelled by adding two **10** hypermultiplets, H_1 and H_2 . The cancellation of the reducible anomalies can be achieved by means of the Green-Schwarz mechanism [7].

For these hypermultiplets we have to define parities with respect to the discrete symmetries,

$$P_I H(x, -y, -z) = \eta_I H(x, y, z) , \quad (8)$$

$$P_{PS} H(x, -y + \pi R_5/2, -z) = \eta_{PS} H(x, y + \pi R_5/2, z) , \quad (9)$$

$$P_{GG} H(x, -y, -z + \pi R_6/2) = \eta_{GG} H(x, y, z + \pi R_6/2) , \quad (10)$$

SO(10)	10			
G_{PS}	(1, 2, 2)	(1, 2, 2)	(6, 1, 1)	(6, 1, 1)
G_{GG}	$\mathbf{5}^*_{-2}$	$\mathbf{5}_{+2}$	$\mathbf{5}^*_{-2}$	$\mathbf{5}_{+2}$
	H^c	H	G^c	G
	$Z_2^{PS} \quad Z_2^{GG}$	$Z_2^{PS} \quad Z_2^{GG}$	$Z_2^{PS} \quad Z_2^{GG}$	$Z_2^{PS} \quad Z_2^{GG}$
H_1	+ +	+ -	- +	- -
H_2	+ -	+ +	- -	- +
H_3	- +	- -	+ +	+ -

Table 1: Parity assignment for the bulk **10** hypermultiplets. $H_1^c = H_d$ and $H_2 = H_u$.

with $\eta_i = \pm 1$ ($i = I, PS, GG$). All hypermultiplets split under the extended 6d supersymmetry into two N=1 4d chiral multiplets, $H = (H, H')$. They have the 6d superpotential interactions [24],

$$S_6 = \int d^6x \left(\int d^2\Theta H'(\partial + \sqrt{2}g\Sigma)H + h.c. \right), \quad (11)$$

where $\partial = \partial_5 - i\partial_6$ and g is the 6d gauge coupling. Invariance of the action then requires that the parities of H and H' are opposite. In the following we denote by η_i the parities of the first 4d chiral multiplet, and we choose $\eta_I = +1$.

The discrete symmetry Z_{PS} implies automatically a splitting between the SU(2) doublets and the SU(3) triplets contained in the **10**-plets. The choice $\eta_{PS} = +1$ leads to massless SU(2) doublets and massive colour triplets (cf. table 1). Choosing further $\eta_{GG} = +1$ for H_1 and $\eta_{GG} = -1$ for H_2 , selects the doublet H^c from the SU(5) $\mathbf{5}^*$ -plet contained in H_1 , and the doublet H from the SU(5) $\mathbf{5}$ -plet of H_2 (cf. table 1). The doublets H^c and H have the quantum numbers of the Higgs fields H_d and H_u in the supersymmetric standard model.

Quarks and leptons can be incorporated by adding **16**-plets, additional **10**-plets etc. in the bulk and on the fixpoints. Without any constraint on the multiplicity of these fields there are many possibilities [20, 22]. It is remarkable that for the complex structure Ω given in eq. (1), the requirement of SO(10) anomaly cancellations in the bulk determines the distribution of multiplets uniquely. As already mentioned, two **10**'s are required to cancel the anomaly of the **45** vector multiplet. Hence, the **(10, 3*)** have to be bulk fields. The anomaly of the third **10**-plet can only be cancelled by the **(16*, 1)**, which leaves **(16, 3)** as brane fields. Note, that in general a bulk field contains two 4-dimensional $N = 1$ chiral multiplets which transform as complex conjugates of each other with respect to the gauge group. Hence, in our SO(10) model in 6 dimensions the content of

SO(10)	16*			
G_{PS}	$(4^*, 2, 1)$	$(4^*, 2, 1)$	$(4, 1, 2)$	$(4, 1, 2)$
G_{GG}	10^*_{+1}	5_{-3}	10^*_{+1}	$5_{-3}, 1_{+5}$
	Q^c	L^c	U^c, E^c	D, N
	$Z_2^{PS} \quad Z_2^{GG}$	$Z_2^{PS} \quad Z_2^{GG}$	$Z_2^{PS} \quad Z_2^{GG}$	$Z_2^{PS} \quad Z_2^{GG}$
Φ^c	$- \quad -$	$- \quad +$	$+ \quad -$	$+ \quad +$

Table 2: Parity assignment for the bulk **16*** hypermultiplet.

4d chiral multiplets is larger than the complex structure (1).

We now have to choose the parities for H_3 , the third **10**-plet, and for Φ^c , the **16***-plet. The presence of Φ^c offers the possibility to break $U(1)_{B-L}$ spontaneously by a vacuum expectation value of its $SU(5)$ singlet component N (table 2). To have N as zero mode fixes the parities of Φ^c to be $\eta_{PS} = \eta_{GG} = -1$. Then an additional colour triplet, D , appears as zero mode. As we shall see, D can acquire a Dirac mass together with another colour triplet, G^c , which can be chosen as zero mode of the third **10**-plet. The corresponding parities of H_3 are $\eta_{PS} = -1$, $\eta_{GG} = +1$. The parities of all components of H_i and Φ^c are listed in tables 1 and 2.

Knowing the parities of all fields we can now discuss the superpotential. We consider the three brane fields **16**, ψ_i , on the $SO(10)$ symmetric fixpoint O . To restrict the number of terms we require R-invariance and an additional global $U(1)_{\tilde{X}}$ symmetry (cf. table 3). The most general superpotential up to quartic terms is then given by,

$$\begin{aligned}
W_4 = & h_d \psi \psi H_1 + h_u \psi \psi H_2 + \frac{h_N}{M_*} \psi \psi \Phi^c \Phi^c \\
& + M_1 H_1 H_3 + M_2 H_2 H_3 + \lambda_1 \Phi^c \Phi^c H_3,
\end{aligned} \tag{12}$$

where we choose $M_* > 1/R_{5,6}$ to be the cutoff of the 6d theory, and the bulk fields have been properly normalised. The first three terms are familiar from ordinary $SO(10)$ GUTs whereas the last three terms are additional couplings among bulk fields.

It is instructive to consider in eq. (12) just the zero mode interactions and their

	H_1	H_2	ψ_i	Φ^c	H_3
R	0	0	1	0	2
$\tilde{X}$	$-2a$	$-2a$	a	$-a$	$2a$

Table 3: Charge assignments for the symmetries $U(1)_R$ and $U(1)_{\tilde{X}}$.

couplings to a single heavy field in the bilinear and cubic terms. In standard notation, with $\psi = (q, u^c, e^c, l, d^c, n^c)$, one obtains,

$$\begin{aligned}
W_4 = & h_d(qd^c + le^c)H_1^c + h_u(qu^c + ln^c)H_2 \\
& + \frac{h_N}{M_*}(n^c N)^2 + \frac{h'_N}{M_*}(d^c D)^2 + \frac{h''_N}{M_*}d^c D n^c N \\
& + \lambda_1 D N G_3^c + M_1 H_1^c H_3 + M_2 H_2 H_3^c \\
& + h_d(qu^c + ln^c)H_1 + h_d(qq + u^c e^c + d^c n^c)G_1 + h_d(ql + u^c d^c)G_1^c \\
& + h_u(qd^c + le^c)H_2^c + h_u(qq + u^c e^c + d^c n^c)G_2 + h_u(ql + u^c d^c)G_2^c ; \quad (13)
\end{aligned}$$

here the three couplings h_N , h'_N and h''_N correspond to the three SO(10) invariants which can be formed from $\psi\psi\Phi^c\Phi^c$.

Vacuum expectation values $v_1 = \langle H_1^c \rangle$, $v_2 = \langle H_2 \rangle$ and $v_N = \langle N \rangle$ yield the mass terms,

$$W_m = m_u u u^c + m_\nu n n^c + m_d d d^c + m_e e e^c + m_N n^c n^c , \quad (14)$$

with mass matrices $m_u = m_\nu = h_u v_2$, $m_d = m_e = h_d v_1$ and $m_N = h_N v_N^2 / M_*$. With $v_{1,2} \ll v_N$ this gives, to first approximation, a good description of the observed properties of quarks and leptons. Since there are two Higgs doublets, H_1 and H_2 , the unwanted relation of minimal SO(10) models, $m_u = m_d$, is avoided. The evidence for small neutrino masses, together with the seesaw mechanism [25], suggests that $U(1)_{B-L}$ is broken near the unification scale, i.e. $v_N \sim 1/R_{5,6}$, which implies large Majorana masses [26]. This also leads to a large mass term, $m_T D G_3^c$, for the colour triplet zero modes of Φ^c and H_3 , with $m_T = \lambda_1 v_N$. The vacuum expectation value $\langle N \rangle = v_N$ breaks $U(1)_{\tilde{X}} \times U(1)_X$ to the diagonal global subgroup, leaving $U(1)_R$ unbroken. As a consequence, integrating out the heavy fields $G_{1,2}$, $G_{1,2}^c$ does not lead to a dimension-5 operator for proton decay. For the same reason, integrating out H_3 and H_3^c does not generate a μ -term $H_1^c H_2$. Finally, the electroweak scale $v_{1,2}$ may be induced together with the μ -term by supersymmetry breaking. The resulting low energy effective theory is just the supersymmetric standard model with right-handed neutrinos.

What determines the vacuum expectation value $\langle N \rangle$ which breaks $U(1)_X$ and therefore $U(1)_{B-L}$? Note, that these $U(1)$ symmetries are anomalous, since the zero modes of Φ^c and H_3 lead to 4d anomalies $SU(3)^2 \times U(1)_X$ and $U(1)_X^3$. In general, these can be compensated by a Chern-Simons term¹. One may then hope to break $U(1)_X$ spontaneously by means of a Fayet-Iliopoulos term on the Georgi-Glashow fixpoint O_{GG} . This will be discussed in more detail elsewhere [27].

Alternatively, one can avoid the occurrence of anomalies by ‘partial doubling’, familiar from supersymmetric σ -models where one associates an entire chiral multiplet with

¹For recent discussions in 5-dimensional theories and references, see [28, 29]

SO(10)	10			
G_{PS}	(1, 2, 2)	(1, 2, 2)	(6, 1, 1)	(6, 1, 1)
G_{GG}	$\mathbf{5}^*_{-2}$	$\mathbf{5}_{+2}$	$\mathbf{5}^*_{-2}$	$\mathbf{5}_{+2}$
	H^c $Z_2^{PS} \quad Z_2^{GG}$	H $Z_2^{PS} \quad Z_2^{GG}$	G^c $Z_2^{PS} \quad Z_2^{GG}$	G $Z_2^{PS} \quad Z_2^{GG}$
H_4	— —	— +	+ —	+ +

SO(10)	16			
G_{PS}	(4, 2, 1)	(4, 2, 1)	(4*, 1, 2)	(4*, 1, 2)
G_{GG}	$\mathbf{10}_{-1}$	$\mathbf{5}^*_{+3}$	$\mathbf{10}_{-1}$	$\mathbf{5}^*_{+3}, \mathbf{1}_{-5}$
	Q $Z_2^{PS} \quad Z_2^{GG}$	L $Z_2^{PS} \quad Z_2^{GG}$	U, E $Z_2^{PS} \quad Z_2^{GG}$	D^c, N^c $Z_2^{PS} \quad Z_2^{GG}$
Φ	— —	— +	+ —	+ +

Table 4: Parity assignment for the bulk **10** and **16** hypermultiplets.

some degrees of freedom of the coset space G/H (cf. [30]). Let us then add two bulk fields, a **16**-plet, Φ , and a fourth **10**-plet, H_4 , which have no irreducible 6d anomaly. The choice of parities $\eta_{PS} = -1$ and $\eta_{GG} = -1$ for both fields leads to the zero modes D^c , N^c and G (cf. table 4). The zero modes of all bulk fields now form a real, and therefore anomaly free, representation of $G_{SM'}$. The additional superpotential terms on the SO(10) brane are,

$$\begin{aligned}
W'_4 = & M^2 S + \lambda_2 S \Phi \Phi^c + \lambda_3 \Phi \Phi H_4 \\
& + \frac{\lambda_4}{M_*} S \Phi \Phi H_1 + \frac{\lambda_5}{M_*} S \Phi \Phi H_2 + \frac{\lambda_6}{M_*} \Phi \Phi^c H_1 H_3 + \frac{\lambda_7}{M_*} \Phi \Phi^c H_2 H_3, \quad (15)
\end{aligned}$$

where we have also included one of the SO(10) singlets, S . The R - and $\tilde{X}$ -charges of the additional fields are $R_\Phi = 0$, $R_{H_4} = R_S = 2$, $\tilde{X}_\Phi = a$, $\tilde{X}_{H_4} = -2a$, $\tilde{X}_S = 0$. Without the singlet field S , there is a bulk D- and F-flat direction, $\langle N \rangle = \langle N^c \rangle = v_N$ [27]. The couplings of S on the brane lift this degeneracy, and one has $v_N = \langle N \rangle = \langle N^c \rangle = \sqrt{-M^2/\lambda_2}$. The cubic term in (15) gives mass to two zero modes G and D^c , $N - N^c$ makes to the $U(1)_X$ vector multiplet massive, and $N + N^c$ has a common mass term with S . Hence, one obtains the same low energy theory as in the anomalous model discussed above.

We have shown that complex structures of coset spaces $E_8/\text{SO}(10) \times H_F$ can provide the starting point of a full grand unified model in the context of supersymmetric SO(10)

theories in 6 dimensions with orbifold compactification. Open questions concern the breaking of supersymmetry, the origin of the brane superpotential and, in particular, the possible connection to theories in 10 dimensions with E_8 symmetry.

We would like to thank A. Hebecker, H. P. Nilles, M. Olechowski and C. Scrucca for helpful discussions.

References

- [1] H. Georgi, S. L. Glashow, Phys. Rev. Lett. **32** (1974) 438
- [2] H. Georgi, Particles and Fields 1974, ed. C. E. Carlson (AIP, NY, 1975) p. 575;
H. Fritzsch, P. Minkowski, Ann. of Phys. **93** (1975) 193
- [3] J. C. Pati, A. Salam, Phys. Rev. **D 10** (1974) 275
- [4] F. Gürsey, P. Ramond, P. Sikivie, Phys. Lett. **B 60** (1976) 177
- [5] N. S. Baaklini, Phys. Lett. **B 91** (1980) 376
- [6] I. Bars, M. Günaydin, Phys. Rev. Lett. **45** (1980) 859
- [7] M. B. Green, J. H. Schwarz, Phys. Lett. **B 149** (1984) 117
- [8] P. Candelas, G. T. Horowitz, A. Strominger, E. Witten, Nucl. Phys. **B 258** (1985) 46
- [9] L. J. Dixon, J. A. Harvey, C. Vafa, E. Witten, Nucl. Phys. **B 261** (1985) 678
- [10] W. Buchmüller, S. T. Love, R. D. Peccei, T. Yanagida, Phys. Lett. **B 115** (1982) 233
- [11] C. L. Ong, Phys. Rev. **D 27** (1983) 3044; *ibid.* Phys. Rev. **D 31** (1985) 3271
- [12] S. Irié and Y. Yasui, Z. Phys. **C29** (1985) 123
- [13] W. Buchmüller, O. Napoly, Phys. Lett. **B 163** (1985) 161
- [14] K. Itoh, T. Kugo, H. Kunitomo, Progr. Theor. Phys. **75** (1986) 386
- [15] Y. Kawamura, Progr. Theor. Phys. **103** (2000) 613; *ibid.* **105** (2001) 999
- [16] G. Altarelli, F. Feruglio, Phys. Lett. **B 511** (2001) 257

- [17] L. J. Hall, Y. Nomura, Phys. Rev. **D 64** (2001) 055003
- [18] A. Hebecker, J. March-Russell, Nucl. Phys. **B 613** (2001) 3
- [19] T. Asaka, W. Buchmüller, L. Covi, Phys. Lett. **B 523** (2001) 199
- [20] L. J. Hall, Y. Nomura, T. Okui, D. R. Smith, Phys. Rev. **D 65** (2002) 035008
- [21] T. Watari, T. Yanagida, hep-ph/0201086
- [22] N. Haba, T. Kondo, Y. Shimizu, hep-ph/0112132; hep-ph/0202191
- [23] A. Hebecker, J. March-Russell, Nucl. Phys. **B 625** (2002) 128
- [24] N. Arkani-Hamed, T. Gregoire, J. Wacker, JHEP **0203** (2002) 055
- [25] T. Yanagida, in *Workshop on unified Theories*, KEK report 79-18 (1979) p. 95;
M. Gell-Mann, P. Ramond, R. Slansky, in *Supergravity* (North Holland, Amsterdam, 1979) eds. P. van Nieuwenhuizen, D. Freedman, p. 315
- [26] E. Witten, Phys. Lett. **B 91** (1980) 81
- [27] T. Asaka, W. Buchmüller, L. Covi, in preparation
- [28] R. Barbieri, R. Contino, P. Creminelli, R. Rattazzi, C. A. Scrucca, hep-th/0203039
- [29] S. Groot Nibbelink, H. P. Nilles, N. Olechowski, hep-th/0203055
- [30] W. Buchmüller, W. Lerche, Ann. Phys. NY **175** (1987) 159